arXiv:0908.2268v1 [q-bio.PE] 17 Aug 2009

Minimisation of a Free-Energy-Like Potential for Non-Equilibrium Systems at Steady State

Robert K. Niven

School of Engineering and Information Technology, The University of New South Wales at ADFA, Canberra, ACT, 2600, Australia.

Abstract

This study examines a new formulation of non-equilibrium thermodynamics, which gives a conditional derivation of the "maximum entropy production" (MEP) principle for flow and/or chemical reaction systems at steady state. The analysis uses a dimensionless potential function ϕ_{st} for non-equilibrium systems, analogous to the free energy concept of equilibrium thermodynamics. Spontaneous reductions in ϕ_{st} arise from increases in the "flux entropy" of the system - a measure of the variability of the fluxes - or in the local entropy production; conditionally, depending on the behaviour of the flux entropy, the formulation reduces to the MEP principle. The inferred steady state is also shown to exhibit high variability in its instantaneous fluxes and rates, consistent with the observed behaviour of turbulent fluid flow, heat convection and biological systems; one consequence is the coexistence of energy producers and consumers in ecological systems. The different paths for attaining steady state are also classified.

Key words: MaxEnt, maximum entropy production, thermodynamics, steady state, flow reaction, irreversible, biological system

---

1. Introduction

Since the seminal book "What is Life" by Erwin Schrödinger (1944), scientists have pondered the existence of life and its compatibility with the second law of thermodynamics. Ridiculing the popular notion that the primary purpose of biological metabolism is to extract matter and energy from the environment, he moves to the crux of the issue (p 71):

> "... *a living organism continually increases its entropy ... and thus tends to approach the dangerous state of maximum entropy, which is death. It can only keep aloof from it, i.e. alive, by continually drawing from its environment negative entropy ... What an organism feeds upon is negative entropy.*"

(The argument is qualified in a footnote, to refer to free energy instead of negative entropy.) The topic was taken up in more detail by Prigogine (1967, 1980), who described living organisms – along with heat-transporting convection cells, turbulent fluid flow vortices and oscillatory chemical reactions – as *dissipative structures*, which continually dissipate heat and thus generate and export entropy to the environment. However, Prigogine's main quantitative result, his *minimum entropy production* (MinEP) principle – valid in the linear or Onsager (1931a,b) transport regime – seems diametrically opposed to life (Martyushev et al., 2007), as was recognised by Prigogine (1980, p88) himself. Bacteria in a microcosm, organisms in an ecosystem or humans on a planet do not try to minimise their entropy production, but instead grow, reproduce and consume all available resources as rapidly as possible. More recently, other thermodynamics-inspired perspectives on biological systems have been advanced, including the use of biological measures of entropy and information (e.g. Ayres, 1994); the non-mathematical gradient theory of Schneider & Sagan (2005); and exergy-based treatments of ecological systems and processes (e.g. Jørgensen, 2006).

Email address: `r.niven@adfa.edu.au` (Robert K. Niven)

Over the past 30 years, a new principle has been proposed, the *maximum entropy production* (MEP) principle, which states that a flow system subject to various flows or gradients will tend towards a steady state position of maximum thermodynamic entropy production, $\dot{\sigma}$ (Ozawa et al., 2003; Kleidon & Lorenz, 2005; Martyushev & Seleznev, 2006; Bruers, 2007c). The MEP principle has been successfully applied – in a heuristic sense – to the prediction of steady states of a wide range of systems, including the Earth's climate system (e.g. Paltridge, 1975, 1978; Kleidon, 2004; Kleidon & Lorenz, 2005); thermal (Bénard) convection (Ozawa et al., 2001); mantle convection (Vanyo & Paltridge, 1981; Lorenz, 2001); electrical currents (Županović et al., 2004; Botric et al., 2005; Christen, 2006; Bruers et al., 2007a); plasmas (Christen, 2007a; Yoshida & Mahajan, 2008); crystalline solids (Martyushev & Axelrod, 2003; Christen, 2007b); ecological systems (Meysman & Bruers, 2007) and biochemical processes (Juretić & Županović, 2003; Dewar et al., 2006). The MEP principle therefore offers a new approach for the analysis of biological systems at the cellular, organism, ecosystem and biosphere levels. Most importantly, it is a *quantitative* principle, based on precisely defined, rigorous thermodynamic concepts; it does not rest upon vague, non-mathematical notions such as "order", "disorder", "randomness" or "complexity" often seen in discussions of biological systems.

Several theoretical justifications of the MEP principle have been advanced, including approaches based on path or transition probabilities (Dewar, 2003, 2005; Attard, 2006a,b) and two more generalistic arguments (Županović et al., 2006; Martyushev, 2007). Recently, a rather different derivation was presented to *directly* determine the steady state of a flow system, based on an entropy defined on the set of local instantaneous fluxes and reaction rates through or within each infinitesimal element; this reduces to a local form of the MEP principle in some circumstances (Niven, 2009b). The analysis invokes a generalised potential function (negative Massieu function) obtained from Jaynes' maximum entropy method, somewhat analogous to the free energy concept used in equilibrium thermodynamics, which attains a minimum at steady state. The *aim* of this study is to explore the implications of this derivation in somewhat simpler terms than in Niven (2009b), using terminology adapted from chemical and statistical thermodynamics. In particular, the nature of the inferred steady state of a flow system, and the various means by which it can be attained, are examined in detail. The analysis has important implications for the modelling of flow systems, including the Earth's climatic-biosphere system and all biological systems.

2. The Generalised Free Energy Concept

2.1. Jaynes' MaxEnt

The maximum entropy (MaxEnt) principle of Jaynes (1957, 1963, 2003, see also Tribus, 1961a,b; Kapur & Kesavan, 1992) provides a powerful technique with which to infer the most probable position of a probabilistic system. Consider a system composed of N distinguishable entities allocated to s equiprobable, distinguishable categories (a *multinomial system*: Niven, 2005, 2006, 2007, 2009a; Niven & Grendar, 2009). In the asymptotic limit $N \to \infty$, the most probable position can be obtained by maximising the relative entropy function (the negative of the Kullback-Leibler Kullback & Leibler (1951) function, D):

$$\mathfrak{H} = -D = -\sum_{i=1}^{s} p_i \ln \frac{p_i}{q_i} \tag{1}$$

where p_i is the probability of an entity in the ith category and q_i is the source or "prior" probability of category i. For equiprobable categories, this reduces to the Shannon (1948) entropy function:

$$\mathfrak{H}_{Sh} = -\sum_{i=1}^{s} p_i \ln p_i \tag{2}$$

plus a constant. Eq. (1) (or (2)) is maximised subject to the natural (normalisation) and any moment constraints on the system:

$$\sum_{i=1}^{s} p_i = 1, \tag{3}$$

$$\sum_{i=1}^{s} p_i f_{ri} = \langle f_r \rangle, \quad r = 1, ..., R, \tag{4}$$

where f_{ri} is the value of the ith category of property f_r and $\langle f_r \rangle$ is the expectation (average) of f_{ri}. This yields the most probable (stationary) distribution of the system:

$$\begin{aligned} p_i^* &= q_i \exp\Big(-\lambda_0^* - \sum_{r=1}^{R} \lambda_r f_{ri}\Big) = (Z^*)^{-1} q_i \exp\Big(-\sum_{r=1}^{R} \lambda_r f_{ri}\Big), \\ Z^* &= q_i \exp(\lambda_0^*) = \sum_{i=1}^{s} \exp\Big(-\sum_{r=1}^{R} \lambda_r f_{ri}\Big), \end{aligned} \tag{5}$$

and the maximum entropy position (Jaynes, 1957, 1963, 2003):

$$\mathfrak{H}^* = \lambda_0^* + \sum_{r=1}^{R} \lambda_r \langle f_r \rangle, \tag{6}$$

where λ_r is the rth Lagrangian multiplier, λ_0^* is the "Massieu function" (Massieu, 1869), Z^* is the partition function and an asterisk denotes the stationary position. It is emphasised that the above derivation is generic, and applies to *any* probabilistic system of multinomial form; it need not refer to a thermodynamic system.

2.2. Generalised Heat, Work and Potential Function

We now consider any conserved quantity f_r, for which we adopt the definition:

$$d\langle f_r \rangle = \delta W_r + \delta Q_r \tag{7}$$

where the path differentials $\delta W_r = \sum_{i=1}^{s} p_i^* d f_{ri}$ and $\delta Q_r = \sum_{i=1}^{s} dp_i^* f_{ri}$ can be termed the "generalised work" and "generalised heat" associated respectively with a change in $\langle f_r \rangle$. It can be shown (Jaynes, 1957, 1963, 2003) that:

$$d\mathfrak{H}^* = \sum_{r=1}^{R} \lambda_r \delta Q_r \tag{8}$$

This is a "generalised Clausius equality" (c.f. Clausius, 1865), applicable to all multinomial systems. Substituting (8) into the differential of (6) and rearranging gives:

$$\begin{aligned} d\phi = -d\lambda_0^* &= \sum_{r=1}^{R} \lambda_r \delta W_r + \sum_{r=1}^{R} d\lambda_r \langle f_r \rangle \\ &= -d\mathfrak{H}^* + d\left(\sum_{r=1}^{R} \lambda_r \langle f_r \rangle\right) \end{aligned} \tag{9}$$

We therefore obtain a potential function ϕ (negative Massieu function) which captures all possible changes in the system, due to changes in the entropy $\mathfrak{H}^*$ or in the "constraint set" $\sum_{r=1}^{R} \lambda_r \langle f_r \rangle$. If the multipliers $\{\lambda_r\}$ are constant, $d\phi$ reduces to the multiplier-weighted total generalised work on the system, $\sum_{r=1}^{R} \lambda_r \delta W_r$. We therefore see that ϕ is a dimensionless, weighted, extended version of the free energy function, applicable to *any* probabilistic system of multinomial form (Jaynes, 1957, 1963, 2003; Tribus, 1961a,b).

How should we interpret (9)? Consider some form of "open system", consisting of a defined region or collection of discrete entities in contact with some surroundings (or the rest of the universe). The internal structure of the system may be described by some probability function p_i, giving rise to some relative entropy function $\mathfrak{H}$ for the system (not necessarily the thermodynamic entropy S, but *any* entropy). From a purely probabilistic formulation of the second law (Niven, 2009a,b):

> *"The entropy of the universe, $\mathfrak{H}_{univ}$, however defined, can only increase",*

it is evident that any spontaneous event must result in an increase of the entropy of the system, $\mathfrak{H}^*$, and/or an increase in entropy produced and exported by the system to its surroundings, $\mathfrak{H}_{prod}$. Quantitatively, this can be written as[1]:

$$d\mathfrak{H}_{univ} = d\mathfrak{H}^* + \delta\mathfrak{H}_{prod} \geq 0 \tag{10}$$

However, the only means by which a system can produce and export entropy – thereby increasing $\delta\mathfrak{H}_{prod}$ – is by a reduction in the magnitude of one or more constraints (or multipliers) which govern the system, $\{\langle f_r \rangle\}$ (or $\{\lambda_r\}$). For such a change to give rise to a quantity in (dimensionless) entropy units, we therefore establish that $\delta\mathfrak{H}_{prod} = -d(\sum_{r=1}^{R} \lambda_r \langle f_r \rangle)$. Comparing (9) and (10), we thus see that $d\phi$ expresses, in a negative sense, the change in entropy of the universe. This can be written as:

$$d\phi = -d\mathfrak{H}^* - \delta\mathfrak{H}_{prod} \leq 0 \tag{11}$$

Eqs. (9) and (11) thus provide a mathematical formulation of a generalised second law (with sign reversed), expressing the interplay between changes in the entropy – however defined – of a system, and changes in entropy produced and exported by a system to its surroundings. This is again consistent with the interpretation of ϕ as a dimensionless, weighted, extended version of the free energy concept (Jaynes, 1957, 1963, 2003; Tribus, 1961a,b).

---

[1] Technically, the variation in $\mathfrak{H}_{prod}$ is written with a δ, since it is a "non-property" of the system; however, for a reproducible phenomenon, it will be expressible in terms of other state functions of the system.

3. Applications

3.1. Equilibrium Systems Example

The above discussion is best illustrated by an example from thermodynamics. Whilst a broader free energy concept is considered in Niven (2009b), most readers will be more familiar with the Gibbs (1875-78) free energy function for systems of constant composition:

$$G = -TS^* + U + PV = -TS^* + H, \tag{12}$$

where S^* is the maximum thermodynamic entropy, U is internal energy, V is volume, P is pressure, T is absolute temperature and H is the enthalpy. Eq. (12) can be derived by applying Jaynes' method to an equilibrium thermodynamic system subject to the constraints U and V, wherein p_i is the joint probability that a molecule will occupy a specified energy level and volume element. Eq. (9) then gives the potential function (Jaynes, 1957, 1963; Tribus, 1961a,b; Jaynes, 2003):

$$d\phi_{eq} = -d\mathfrak{H}^*_{eq} + d(\lambda_U U + \lambda_V V) \tag{13}$$

Recognising $S^* = k\mathfrak{H}^*_{eq}$, $\lambda_U = 1/kT$ and $\lambda_V = P/kT$, where k is Boltzmann's constant, gives:

$$k d\phi_{eq} = d\Big(\frac{G}{T}\Big) = -dS^* + d\Big(\frac{H}{T}\Big) \leq 0 \tag{14}$$

equivalent to (12). This form reveals the true meaning of the Gibbs free energy concept: it expresses – in a negative sense – the interplay between the change in entropy within the system, dS^*, and the change in entropy exported by the system, $-d(H/T)$, due to transfers of heat (Planck, 1922, 1932; Fermi, 1936; Strong & Halliwell, 1970; Craig, 1988). From the (classical) second law, their sum must be positive, and so a system will spontaneously approach a position of minimum G/T (for constant T, it will approach minimum G). From (11), we can rewrite (14) as (Niven, 2009b):

$$k d\phi_{eq} = d\Big(\frac{G}{T}\Big) = -dS^* - \delta\sigma \leq 0 \tag{15}$$

where $\delta\sigma = k\delta\mathfrak{H}_{prod}$ is the increment of *thermodynamic entropy produced and exported* by the system to its surroundings.

3.2. Flow System Example

Now consider a second example, of an infinitesimal fluid element in a control volume of a flowing fluid, subject to local mean values of the heat flux $\boldsymbol{j}_Q$, diffusive mass fluxes $\boldsymbol{j}_c$ of each chemical species c, stress tensor $\boldsymbol{\tau}$ and chemical reaction rates $\hat{\dot{\xi}}_d$ of each reaction d, plus the natural constraint (3). This model encompasses all biological and ecological systems. Such a system can be analysed by Jaynes' method using the local flux relative entropy (Niven, 2009b):

$$\mathfrak{H}_{st} = -\sum_{\boldsymbol{I}} \pi_{\boldsymbol{I}} \ln \frac{\pi_{\boldsymbol{I}}}{\gamma_{\boldsymbol{I}}} \tag{16}$$

where $\pi_{\boldsymbol{I}}$ is the joint probability that the fluid element experiences a set of instantaneous local fluxes of heat, species c, momentum and rates of chemical reactions d, and $\gamma_{\boldsymbol{I}}$ is the joint prior probability. For this system, it can be shown that (9) gives the increment in the local potential function (Niven, 2009b):

$$d\phi_{st} = -d\mathfrak{H}^*_{st} - \frac{\theta\mathcal{V}}{k}\delta\hat{\dot{\sigma}} \leq 0 \tag{17}$$

where $\mathfrak{H}^*_{st}$ is the local flux entropy at steady state, θ and $\mathcal{V}$ are characteristic time and volume scales of the system, and $\hat{\dot{\sigma}}$ is the local *thermodynamic entropy production* of the element per unit volume:

$$\begin{aligned} \hat{\dot{\sigma}} = \boldsymbol{j}_Q \cdot \boldsymbol{\nabla}\Big(\frac{1}{T}\Big) - \sum_c \boldsymbol{j}_c \cdot \Big[\boldsymbol{\nabla}\Big(\frac{\mu_c}{M_c T}\Big) - \frac{\boldsymbol{g}_c}{T}\Big] \\ -\boldsymbol{\tau} : \boldsymbol{\nabla}\Big(\frac{\boldsymbol{v}}{T}\Big)^{\top} - \sum_d \hat{\dot{\xi}}_d \frac{A_d}{T} \end{aligned} \tag{18}$$

in which μ_c, M_c and $\boldsymbol{g}_c$ are the molar chemical potential, molar mass and specific body force on species c, $\boldsymbol{v}$ is the mass-average fluid velocity and A_d is the molar chemical affinity of reaction

d, with $A_d < 0$ indicating spontaneous forwards reaction. Eq. (18) can be summarised in the form:

$$\hat{\dot{\sigma}} = \sum_X j_X F_X \tag{19}$$

where j_X is the local mean flux or mean reaction rate of quantity or species X, and F_X is the corresponding local "thermodynamic force" (gradient or affinity term). Not coincidentally, (17) has the same form as (11), expressing (with sign reversed) the sum of changes in the flux entropy within the element plus its export out of the system. A flow element will therefore try to approach a steady state position of minimum ϕ_{st}, for the same reason that an equilibrium thermodynamic system tries to approach an equilibrium position of minimum ϕ_{eq} (minimum G/T).

Comparing (14) and (17), we see that the entropy production $\hat{\dot{\sigma}}$ within a fluid element of a steady state system plays a similar role (with change of sign and units) to the enthalpy function H in equilibrium systems. This is an important insight, which has perhaps been hindered by the lack of popular understanding of the free energy concept (see Strong & Halliwell, 1970; Craig, 1988). The common feature is that H and $\hat{\dot{\sigma}}$ both serve as (modified) measures of the export of entropy – however defined – by a system to its surroundings. Many previous authors have erred in considering $\hat{\dot{\sigma}}$ to be the non-equilibrium analogue of S^*; whilst this may seem reasonable at first glance, it is not correct.

4. Implications

4.1. Meaning of the Flux Entropy

To understand (17), it is necessary to appreciate the meaning of the flux entropy $\mathfrak{H}^*_{st}$. To do this, we need to consider the mathematical properties of the Shannon entropy (2) (e.g. Kapur & Kesavan, 1992, chap. 2). In essence, $\mathfrak{H}$ indicates the spread of the distribution p_i amongst its categories i; the thermodynamic entropy S^* therefore reflects the spread of the equilibrium distribution p_i^* over energy levels and volume elements, with low S^* indicating a narrow distribution and high S^* a broad one. In the same way, $\mathfrak{H}^*_{st}$ reflects the spread of the steady state probability $\pi^*_{\boldsymbol{I}}$ over the set of instantaneous local fluxes and reaction rates. This is illustrated by the schematic plots in Figures 1a-c, for a univariate parameter $\boldsymbol{I} = I$ (e.g., a single flux or reaction rate of quantity X). All three plots have the same mean flux or rate j_X – represented schematically by a fixed "mean category" I_0 – but the variance, and therefore the flux entropy, increases to the right.

The plots reveal an additional, extraordinary feature of flow systems. In equilibrium systems, the categories (e.g. energy levels) are generally taken to start from a "zero" or reference level, for which the value of the index is unimportant. In contrast, flow systems have no such minimum, since we must allow for positive and negative flux or rate levels $I = 0, \pm 1, \pm 2, ...$ In consequence, as $\mathfrak{H}^*_{st}$ increases, the system is more likely to access its states of reverse flow or reverse chemical reaction $I < 0$, even if the mean value j_X is high (see Figures 1b-c). In other words, a high flux entropy is associated with *greater variability* in the fluxes and rates, which therefore implies *oscillatory* or *chaotic* processes. We immediately see the connection between steady states of high $\mathfrak{H}^*_{st}$ and the defining features of many "far from equilibrium" systems, such as fluid turbulence, heat-induced convection cells, nonlinear diffusion phenomena and oscillatory chemical reactions (Prigogine, 1967, 1980); indeed, the latter are prevalent in biochemical processes such as nutrient degradation processes (Meysman & Bruers, 2007) and the photosynthesis cycle (Juretić & Županović, 2003; Dewar et al., 2006).

The importance of above analysis can be illustrated by its application to the species population structure within an ecosystem. Consider a small element of a (rudimentary) ecosystem of s species, identified only by their energy usage, such that each organism of species i has the energy consumption ϵ_i, whilst the system has mean energy consumption $\langle E \rangle$. This model dramatically simplifies the MaxEnt ecosystem model given by Dewar & Porté (2008). To infer the steady state population distribution π_i^*, we maximise the relative entropy $\mathfrak{H}_{ecol} = -\sum_{i=1}^{s} \pi_i \ln \pi_i / \gamma_i$ (16), subject to known prior probabil-


$\mathfrak{H}^*_{st1}$
π^*_I
0
0
I_0
I
(a)
$\mathfrak{H}^*_{st2} > \mathfrak{H}^*_{st1}$
π^*_I
0
0
I_0
I
(b)
$\mathfrak{H}^*_{st3} > \mathfrak{H}^*_{st2}$
π^*_I
0
0
I_0
I
(c)

Figure 1: Effect of increasing $\mathfrak{H}^*_{st}$ on a univariate steady-state distribution π^*_I (schematic only).

ities γ_i and constraints $\sum_{i=1}^{s} \pi_i = 1$ and $\sum_{i=1}^{s} \pi_i \epsilon_i = \langle E \rangle$, giving:

$$\begin{aligned} \pi_i^* &= \mathcal{Z}^{-1} \gamma_i \exp(-\zeta_E \epsilon_i) \\ \mathcal{Z} = e^{\zeta_0} &= \sum_{i=1}^{s} \gamma_i \exp(-\zeta_E \epsilon_i) \end{aligned} \tag{20}$$

where ζ_0 and ζ_E are, respectively, the Lagrangian multipliers for the two constraints, and $\mathcal{Z}$ is the partition function. The analysis (17)-(19) then follows from (20), with the energetic multiplier identified as $\zeta_E \propto F_E = \boldsymbol{\nabla} T^{-1}$. Although (20) has the appearance of a Boltzmann distribution akin to that of chemical thermodynamics, in an ecosystem the "energy levels" i are actually "energy consumption levels", which can be positive or negative, corresponding respectively to *net energy consumers* ($i > 0$) and *net energy producers* ($i < 0$). At a high ecological flux entropy $\mathfrak{H}^*_{ecol}$, the "most probable" ecosystem will therefore be forced to contain both energy producers and consumers, rather than just energy consumers. Similarly, in turbulent fluid flow, some energetic structures will be net energy consumers (dissipating energy as heat), whilst others will be net energy producers (transferring energy from its incoming source to the consumers). We therefore recover the essence of the ubiquitous "food chain" (or "food web") of ecological systems and the "energy cascade" of turbulent flow systems.

4.2. *Classification of Spontaneous Processes*

We can now return to the equilibrium (14) and steady state (17) potential functions. For simplicity, we first confine the discussion to processes with monotonic changes in the entropy S^* and entropy produced $\sigma = -H/T$. We see that in equilibrium systems, the path towards equilibrium $d\phi_{eq} \leq 0$ will depend on the relative changes in S^* and σ, leading to three possible scenarios for a spontaneous process, as listed in Table 1. In Case E1, the process is driven by changes in both entropy terms, whilst in Cases E2 and E3, a reduction in one entropy is "paid for" by a greater and opposite gain in the other. In all cases, since the entropy changes are monotonic, the equilibrium position (minimum ϕ_{eq} = minimum G/T) must coincide with extrema (a minimum or maximum) in both S^* and σ, as set out in the Table.

Similarly, from (17), in a flow system subject to monotonic changes in $\mathfrak{H}^*_{st}$ and $\hat{\sigma}$, each increment towards steady state $d\phi_{st} \leq 0$ could be achieved by one of the three cases listed in Table 1. The corresponding extrema at steady state are also listed. As shown, Cases S1 and S3 are consistent with a position of *maximum entropy production* (MEP). Case S2, on the other hand, involves convergence towards a position of *minimum entropy production* (MinEP). The three cases therefore encompass the two major (seemingly contradictory) principles of non-equilibrium thermodynamics (Prigogine, 1967, 1980; Martyushev et al., 2007).

We further note that if passage to equilibrium or steady state is not monotonic, many more scenarios are possible. In thermodynamics, this is handled by considering only the net change in Gibbs free energy $\Delta G = -T\Delta S^* + \Delta H$ at constant T and P. In light of (14), this is more appropriately written as:

$$k\Delta\phi_{eq} = \Delta\Big(\frac{G}{T}\Big) = -\Delta S^* + \Delta\Big(\frac{H}{T}\Big) \leq 0 \tag{21}$$

This rests on the fact that G, S^*, U, V and H are state functions, so we can disregard the path taken by the system. Although such systems could follow any of the paths E1-E3 in Table 1

Table 1: List of possible spontaneous processes in equilibrium and steady state systems, for monotonically varying parameters (terminology from ancient Greek: *exo-*, external; *endo-*, internal; *pan-*, everywhere; *tropos*, transformation (Clausius, 1865) and *-genic*, generating or producing).

Case	Conditions		Entropic Driving Force	Label	Extrema at Stationary Position	
Equilibrium systems ($kd\phi_{eq} = d(G/T) \leq 0$)						
E1	$dS^* \geq 0$,	$\delta\sigma \geq 0$	universal	*panentropic*	max S^*,	max σ (= min H/T)
E2	$dS^* \geq \lvert\delta\sigma\rvert \geq 0$,	$\delta\sigma \leq 0$	internal dominant	*endoentropic*	max S^*,	min σ (= max H/T)
E3	$dS^* \leq 0$,	$\delta\sigma \geq \lvert dS^*\rvert \geq 0$	external dominant	*exoentropic*	min S^*,	max σ (= min H/T)
Steady state systems ($Kd\phi_{st} \leq 0$ with $K = k/\theta\mathcal{V}$)						
S1	$d\mathfrak{H}^*_{st} \geq 0$,	$\delta\hat{\dot{\sigma}} \geq 0$	universal	*panentropogenic*	max $\mathfrak{H}^*_{st}$,	max $\hat{\dot{\sigma}}$
S2	$Kd\mathfrak{H}^*_{st} \geq \lvert\delta\hat{\dot{\sigma}}\rvert \geq 0$,	$\delta\hat{\dot{\sigma}} \leq 0$	internal dominant	*endoentropogenic*	max $\mathfrak{H}^*_{st}$,	min $\hat{\dot{\sigma}}$
S3	$d\mathfrak{H}^*_{st} \leq 0$,	$\delta\hat{\dot{\sigma}} \geq \lvert Kd\mathfrak{H}^*_{st}\rvert \geq 0$	external dominant	*exoentropogenic*	min $\mathfrak{H}^*_{st}$,	max $\hat{\dot{\sigma}}$



Figure 2: Possible responses of a flow system to an increasing force F_X or mean flux j_X (increasing I_0): (a)-(b)-(c) constant $\mathfrak{H}^*_{st}$ or (a)-(d)-(e) increasing $\mathfrak{H}^*_{st}$.

during different stages of the process – or even temporarily deviate from $d(G/T) \leq 0$ to overcome an activation energy barrier – they must approach a position of minimum G/T, leading to a net change $\Delta(G/T) \leq 0$. The system can still be said to follow one of Cases E1-E3, but now only in a net sense (using Δ's rather than *d*'s). In Case E3, for example, we can still speak of the system tending towards a position of minimum S^* and maximum σ (= minimum H/T), provided this is understood to refer to their net changes rather than the path taken by the system.

In a similar vein, if an unsteady flow system is not restricted to purely monotonic changes, it must still approach a steady state position of minimum $\mathfrak{H}^*_{st}$, and thus undergo the net change $\Delta\mathfrak{H}^*_{st} \leq 0$. Presuming that $\mathfrak{H}^*_{st}$ and $\hat{\dot{\sigma}}$ can be considered as state functions, the system can still be identified as following – now in a net sense – one of the three Cases S1-S3 in Table 1.

Can we infer anything more about flow systems? Indeed, we can. Consider a flow element which experiences a gradual increase in the local thermodynamic force F_X conjugate to the mean local flux or rate j_X. Such an element may undergo two types of changes: (i) an increase in j_X without any corresponding increase (or even a decrease) in $\mathfrak{H}^*_{st}$, illustrated in Figures 2a-b-c; and (ii) increases in both j_X and $\mathfrak{H}^*_{st}$, illustrated in Figures 2a-d-e. From (18), both scenarios involve identical increases in the entropy production, $\Delta\hat{\dot{\sigma}} = \Delta(j_X F_X)$. Which is more likely? From our knowledge of flow systems, the first scenario seems less credible, since it requires the fluxes to remain within a narrow range of instantaneous values at all times, even though the driving force has increased. A decrease in $\mathfrak{H}^*_{st}$ seems even more unlikely. The second scenario permits greater variability (fluctuations) of the fluxes, consistent with the formation of a non-linear mechanism to enable greater transport or production of X. A similar argument applies if the flux, rather than the gradient, is the control variable. Although this is not a proof, it does lend support to the argument (Niven, 2009b) that fluid elements tend to undergo concurrent increases in $\mathfrak{H}^*_{st}$ and $\hat{\dot{\sigma}}$, and thus converge to steady state by a (net) *panentropogenic* process. In such cases, the steady state position can be determined by the (net) MEP principle, without concern over contrary effects due to decreases in $\mathfrak{H}^*_{st}$.

5. The MEP "Heuristic"

We now turn to a discussion of current practice in the application of the MEP principle to flow or chemical reaction systems, including biological systems. From the pioneering works of Paltridge (1975, 1978) and three decades of further experience (e.g. Ozawa et al., 2001, 2003; Juretić & Županović, 2003; Kleidon, 2004; Kleidon & Lorenz, 2005; Dewar et al., 2006; Martyushev & Seleznev, 2006; Bruers, 2007c; Meysman & Bruers, 2007), this has evolved into a set of practices which can be termed the "MEP heuristic":

(i) Divide the control volume into very large subdomains (or even consider the entire domain);

(ii) Set up the set of mass, chemical species, energy, momentum and/or charge balance equations for the system, based on the bulk flow rates between subdomains, using linear (Onsager-like) transport equations with adjustable, whole-subdomain transport coefficient(s), and chemical reaction rate equations with adjustable first-order rate constant(s);

(iii) Calculate the thermodynamic entropy production of the system, as a function of the adjustable parameter(s);

(iv) The inferred steady state of the system is given by the position of maximum entropy production with respect to the adjustable parameter(s).

How does this heuristic work? In effect, it selects the highest allowable entropy production consistent with the set of allowable bulk net fluxes $J_{X,\Gamma}$ and bulk thermodynamic forces $F_{X,\Gamma}$ in and between subdomains Γ of the system:

$$\text{MEP\_ Heuristic} = \max_{\boldsymbol{\Omega}} \left(\sum_{\Gamma} \sum_{X} J_{X,\Gamma}(\boldsymbol{\Omega})\, F_{X,\Gamma}(\boldsymbol{\Omega}) \right) \tag{22}$$

where the bulk fluxes and/or forces are functions of the set of subdomain-wide adjustable parameters $\boldsymbol{\Omega} = \{\Omega\}$. It must be recognised, however, that the adjustable parameters are secondary variables, which do not represent fundamental physical processes. The true maximum must therefore be given by a "system maximum entropy production" (SMEP) principle:

$$\text{SMEP} = \max\left(\iiint_{CV} \hat{\dot{\sigma}}(V)\, dV \right) \tag{23}$$

where the maximum is taken with respect to the instantaneous fluxes $j_{X,\boldsymbol{I}}$, conditioned by the constraints on the system, and the integral is calculated over the control volume. The MEP heuristic therefore makes the assumption that (22) and (23) are equivalent, which is correct if and only if there exists a set of local physical mechanisms by which the maximum in (22) can be physically realised. Using the terminology of MEP practitioners, the MEP heuristic (22) must be considered to apply only to "many-degree-of-freedom" systems (e.g. Ozawa et al., 2001, 2003; Juretić & Županović, 2003; Kleidon, 2004; Kleidon & Lorenz, 2005; Dewar et al., 2006; Martyushev & Seleznev, 2006; Bruers, 2007c; Meysman & Bruers, 2007).

In contrast, the analysis herein (§2-4) and in Niven (2009b) gives the optimisation principle:

$$\begin{aligned} \text{Optimum} &= \iiint_{CV} \min(\phi_{st}(V))\, dV \\ &= \iiint_{CV} \max\Big(\mathfrak{H}^*_{st}(V) + \frac{\theta \mathcal{V}\, \hat{\dot{\sigma}}(V)}{k} \Big)\, dV \end{aligned} \tag{24}$$

If the parameters $\mathfrak{H}_{st}^*$ and $\hat{\dot{\sigma}}$ are positively correlated – as argued in §4 – then (24) becomes functionally equivalent to a "local maximum entropy production" (LMEP) principle, which gives for the overall system:

$$\text{LMEP} = \iiint\limits_{CV} (\max \hat{\dot{\sigma}}(V))\, dV \tag{25}$$

This is a much stronger condition than (23). By considerations of integral calculus (Zwillinger, 2003), the two bounds are related by:

$$\max\left(\iiint\limits_{CV} \hat{\dot{\sigma}}(V)\, dV \right) \leq \iiint\limits_{CV} \max \hat{\dot{\sigma}}(V)\, dV \tag{26}$$

since the left hand side could possess regions of $\hat{\dot{\sigma}} < 0$, compensated by other regions of greater $\hat{\dot{\sigma}} > 0$. This, however, runs against an argument used by Prigogine (1967, 1980): how can a system possibly "know" that it can consume entropy in some regions, which will be compensated by greater entropy production in others? Indeed, we could construct a smaller control volume containing only the entropy-consuming elements, which would continuously violate the second law of thermodynamics. It is for this sound reason that the MEP principle must be a local principle, applicable at all volume scales. With the restriction $\hat{\dot{\sigma}}(V) \geq 0$, we see that the two maxima in (26) coincide, and so the MEP heuristic (with its assumption of many degrees of freedom) becomes equivalent to the local formulation.

6. Conclusions

This study examines the meaning and implications of a new formulation of non-equilibrium thermodynamics applicable to flow and/or chemical reaction systems at steady state (Niven, 2009b). This provides a very different, conditional derivation of the "maximum entropy production" (MEP) principle, based on minimisation of a dimensionless, local, free-energy-like potential function ϕ_{st}. The analysis encompasses all biological and ecological systems. Firstly, the basis of the derivation and the meaning of ϕ_{st} are examined. The flux entropy $\mathfrak{H}_{st}^*$ used in the analysis is then shown to represent the "spread" of the distribution of instantaneous fluxes and/or reaction rates through or within the element. Since a flow system can access states of reverse flow or reaction, a high flux entropy is consistent with *higher variability* and thus with *chaotic* or *oscillatory* processes. In this respect, the term "steady state" is therefore something of a misnomer, since it refers only to the constancy of the mean bulk flows and not their temporal and spatial variability. One consequence, examined through a specific example, is the coexistence of energy producers and consumers in ecological systems.

The effects of reinforcement or competition between changes in flux entropy $\mathfrak{H}_{st}^*$ and entropy production $\hat{\dot{\sigma}}$ are then examined and classified. It is argued that in many systems, these two parameters should increase concurrently, enabling the steady state position to be determined by the MEP principle. The "MEP heuristic" used by MEP practitioners is then shown to be consistent with the present local formulation, with the additional assumption that the system has sufficient dynamic degrees of freedom that the MEP state can be physically realised.

Acknowledgments

The author thanks the participants of the MEP workshops hosted by the Max-Planck-Institut für Biogeochemie, Jena, Germany, in 2007 and 2008, for valuable discussions; The University of New South Wales and the above Institute for financial support; and the European Commission for financial support as a Marie Curie Incoming International Fellow (2007-2008) under Framework Programme 6.

References

Attard P. (2006), Statistical mechanical theory for steady state systems. VI. Variational principles, J. Chem. Phys. 125: article 214502.

Attard, P. (2006) Theory for non-equilibrium statistical mechanics, Phys. Chem. Chem. Phys. 8: 3585-3611.

Ayres, R.U. (1994) Information, Entropy and Progress, AIP, Woodbury, NY.

Bénard, H. (1901) Les tourbillons cellulaires dans une nappe liquide transportant de la chaleur par convection en régime permanent, Ann. Chim. Phys. 23: 62-144.

Botrić, S., Županović, P., Juretić, D. (2005) Is the stationary current distribution in a linear planar electric network determined by the principle of maximum entropy production?, Croatica Chemica Acta 78(2): 181-184.

Bruers, S. (2007) Classification and discussion of macroscopic entropy production principles, *arXiv:cond-mat/0604482v3*.

Bruers, S., Maes, C., Netočný, K. (2007) On the validity of entropy production principles for linear electrical circuits, J. Stat. Phys. 129: 725-740.

Christen, T. (2006), Application of the maximum entropy production principle to electrical systems, J. Phys. D: Appl. Phys. 39: 4497-4503.

Christen, T. (2007a), A maximum entropy production model for Teflon ablation by arc radiation, J. Phys. D: Appl. Phys. 40: 5719-5722.

Christen, T. (2007b)Modelling diffusion in nonuniform solids using entropy production rate, J. Phys. D: Appl. Phys. 40: 5723-5726.

Clausius, R. (1865) Über verschiedene für die Anwendung bequeme Formen der Hauptgleichungen der mechanischen WŁrmetheorie, Poggendorfs Annalen 125: 335-400.

Craig, N.C. (1988) Entropy analysis of four familiar processes, J. Chem. Ed. 65(9): 760-764.

Dewar, R.C. (2003) Information theory explanation of the fluctuation theorem, maximum entropy production and self-organized criticality in nonequilibrium stationary states, J. Phys. A: Math. Gen. 36: 631-641.

Dewar, R.C. (2005)Maximum entropy production and the fluctuation theorem, J. Phys. A: Math. Gen. 38: L371-L381.

Dewar, R.C., Juretić, D., Županović, P. (2006)The functional design of the rotary enzyme ATP synthase is consistent with maximum entropy production, Chemical Physics Letters 430: 177-182.

Dewar, R.C., Porté, A. (2008)Statistical mechanics unifies different ecological patterns, J. Theor. Biol. 251: 389-403.

Edelstein-Keshet, E. (1988) Mathematical Models in Biology, SIAM, Philadelphia, PA, USA.

Fermi, E. (1956)Thermodynamics, Dover Publ., NY.

Gibbs, J.W. (1875-78) On the equilibrium of heterogeneous substances, *Trans. Connecticut Acad.* 3 (1875-1876) 108-248; (1877-1878) 343-524.

Jørgensen, S.E. (2006) Eco-Exergy as Sustainability, WIT Press, Southhampton, UK.

Jaynes, E.T. (1957), Information theory and statistical mechanics, Physical Review, 106: 620-630.

Jaynes, E.T. (1963), Information theory and statistical mechanics, *in* Ford, K.W. (ed), Brandeis University Summer Institute, Lectures in Theoretical Physics, Vol. 3: Statistical Physics, Benjamin-Cummings Publ. Co., 181-218.

Jaynes, E.T. (G.L. Bretthorst, ed.) (2003) *Probability Theory: The Logic of Science*, Cambridge U.P., Cambridge.

Juretić, D., Županović, P. (2003) Photosynthetic models with maximum entropy production in irreversible charge transfer steps, Computational Biology and Chemistry 27: 541-553.

Kapur, J.N., Kesevan, H.K. (1992) *Entropy Optimization Principles with Applications*, Academic Press, Inc., Boston, MA.

Kleidon, A. (2004) Beyond Gaia: thermodynamics of life and Earth system functioning, Climatic Change 66: 271-319.

Kleidon, A., Lorenz R.D. (2005) (eds.) Non-equilibrium Thermodynamics and the Production of Entropy: Life, Earth and Beyond, Springer Verlag, Heidelberg.

Kullback, S., Leibler, R.A. (1951) On information and sufficiency, Annals Math. Stat. 22: 79-86.

Lorenz, R.D. (2001) Of course Ganymede and Callisto have oceans: Application of a principle of maximum entropy production to icy satellite convection, Proc. Lunar Planet Sci. Conf., 32: abstract 1160.

Martyushev, L.M. (2007) Do nonequilibrium processes have features in common?, arXiv:0709.0152v1.

Martyushev, L.M., Axelrod, E.G. (2003) From dendrites and S-shaped growth curves to the maximum entropy production principle, JETP Letters 78(8): 476-479.

Martyushev, L.M., Seleznev, V.D. (2006) Maximum entropy production principle in physics, chemistry and biology, Physics Reports 426: 1-45.

Martyushev, L.M., Nazarova, A.S., Seleznev, V.D. (2007) J. Phys. A: Math. Theor. 40: 371-380.

Massieu, M. (1869) Thermodynamique - Sur les fonctions caractéristiques des divers fluides, Comptes Rendus 69: 858-862; 1057-1061.

Meysman, F.J.R., Bruers, S. (2007) A thermodynamic perspective on food webs: Quantifying entropy production within detrital-based ecosystems, J. Theor. Biol. 249: 124-139.

Moran, M.J., Shapiro, H.N. (2006), Fundamentals of Engineering Thermodynamics, 5th ed., John Wiley, NY.

Niven, R.K. (2005), Exact Maxwell-Boltzmann, Bose-Einstein and Fermi-Dirac statistics, Physics Letters A, 342(4): 286-293.

Niven, R.K. (2006), Cost of s-fold decisions in exact Maxwell-Boltzmann, Bose-Einstein and Fermi-Dirac statistics, Physica A, 365(1): 142-149.

Niven, R.K. (2007) Origins of the combinatorial basis of entropy, in Knuth, K.H., Caticha, A., Center, J.L., Giffon, A., Rodrguez, C.C. (eds), MaxEnt07, AIP Conference Proceedings 954, Melville, pp 133-142.

Niven, R.K. & Grendar, M. (2009) Generalized classical, quantum and intermediate statistics and the Polya urn model, Physics Letters A 373: 621-626.

Niven, R.K. (2009a) Combinatorial entropies and statistics, European Physics Journal B 70: 49-63.

Niven, R.K. (2009b) Steady State of a dissipative flow-controlled system and the maximum entropy production principle, Phys. Rev. E, in press; see http://arxiv.org/abs/0902.1568.

Onsager, L. (1931a) Reciprocal relations in irreversible processes I, Phys. Rev. 37: 405-426.

Onsager, L. (1931b) Reciprocal relations in irreversible processes II, Phys. Rev. 38: 2265-2279.

Ozawa, H., Shikokawa, S., Sakuma, H. (2001) Thermodynamics of fluid turbulence: A unified approach to the maximum transport properties, Phys. Rev. E 64: article 026303.

Ozawa, H., Ohmura, A., Lorenz, R.D., Pujol, T. (2003) The second law of thermodynamics and the global climate system: A review of the maximum entropy production principle, Rev. Geophys. 41: article 4.

Paltridge, G.W. (1975) Global dynamics and climate - a system of minimum entropy exchange, Quart. J. Royal Meteorol. Soc. 101: 475-484.

Paltridge, G.W. (1978) The steady-state format of global climate, Quart. J. Royal Meteorol. Soc. 104: 927-945.

Planck, M. (1922) Treatise on Thermodynamics, Engl. transl., 3rd ed., Dover Publications, NY.

Planck, M. (1932) Introduction to Theoretical Physics, Vol. V: Theory of Heat, Engl. transl. H.L. Brose, Macmillan & Co., Ltd.

Prigogine, I. (1967), Introduction to Thermodynamics of Irreversible Processes, 3rd ed., Interscience Publ., NY.

Prigogine, I. (1980), From Being to Becoming: Time and Complexity in the Physical Sciences, W.H. Freeman & Co., San Francisco, 1980.

Schneider, E.D., Sagan, D. (2005) Into the Cool: Energy Flow, Thermodynamics and Life, University of Chicago Press, Chicago.

Schrödinger, E. (1944), What is Life?, Cambridge U.P., Cambridge (see chapter 6 and following Note).

Shannon, C.E. (1948) A mathematical theory of communication, Bell Sys. Tech. J. 27: 379-423; 623-659.

Strong, L.E., Halliwell, H.F. (1970) An alternative to free energy for undergraduate instruction, J. Chem. Ed. 47(5): 347-352.

Tribus, M. (1961a), Information theory as the basis for thermostatics and thermodynamics, J. Appl. Mech., *Trans. ASME*, 28: 1-8.

Tribus, M. (1961b), Thermostatics and Thermodynamics, D. Van Nostrand Co. Inc., Princeton, NJ.

Vanyo, J.P., Paltridge, G.W. (1981) A model for energy dissipation at the mantle-core boundary, Geophys. J. R. Astron. Soc. 66: 677-690.

Yoshida, Z., Mahajan, S.M. (2008) Maximum entropy production in self-organized plasma boundary layer: A thermodynamic discussion about turbulent heat transport, Physics of Plasmas 15: article 032307.

Županović, P., Juretić, D., Botrić, S. (2004)Kirchhoffs loop law and the maximum entropy production principle, Phys. Rev. E 70: article 056108.

Županović, P., Botrić, S., Juretić, D. (2006) Relaxation processes, MaxEnt formalism and Einstein's formula for the probability of fluctuations, Croatica Chemica Acta 79(3): 335-338.

Zwillinger, D. (2003)CRC Standard Mathematical Tables and Formulae, Chapman & Hall / CRC Press, Boca Raton, FL.